

# INDEX TO VOLUME 85 (1999)

## AUTHORS

- Aasen, G.A.: Juvenile delta smelt use of shallow-water and channel habitats in the Sacramento-San Joaquin Estuary, 161-169.
- Barrett, R.H.: see Boroski and Barrett.
- Barrett, R.H.: see Timossi, Garrison, and Barrett.
- Baxter, J.: see Jow, Baxter, and Klingbeil.
- Baxter, R.D.: Status of splittail in California, 28-30.
- Bettaso, R.H. and J.N. Young: Evidence for freshwater spawning by striped mullet and return of the Pacific tenpounder in the lower Colorado River, 75-76.
- Bleich, V.C. and A.M. Pauli: Distribution and intensity of hunting and trapping activity in the East Mojave National Scenic Area, California, 148-160.
- Bleich, V.C. and B.M. Pierce: Expandable and economical long-term collars for juvenile mule deer, 56-62.
- Boroski, B.B. and R.H. Barrett: Movement patterns and survivorship of black-tailed deer migrating across Trinity Reservoir, 63-69.
- Bursey, C.R.: see Goldberg and Bursey.
- Castro-Aguirre, J.L.: see Gonzalez-Acosta, De la Cruz-Agüero, Rodriguez-Sanchez, De la Cruz-Agüero, and Castro-Aguirre.
- Cech, J.J., Jr.: see Magee, Myrick, and Cech.
- Chen, H.-C.: see Liu, Yang, and Chen.
- Childers, J.: see Schaefer and Childers.
- Clark, H.O., Jr.: An animal track casting method using dolomite, sand, and spray adhesive, 138-139.
- Davis, L.H.: Review of: Avian conservation. Research and management, 144-145.
- De la Cruz-Agüero, G.: see Gonzalez-Acosta, Cruz-Agüero, Rodriguez-Sanchez, Cruz-Agüero, and Castro-Aguirre.
- De la Cruz-Agüero, J.: A first Mexican record of the chinook salmon, *Oncorhynchus tshawytscha*, 77-78.
- De la Cruz-Agüero, J.: see Gonzalez-Acosta, Cruz-Agüero, Rodriguez-Sanchez, Cruz-Agüero, and Castro-Aguirre.
- Erickson, R.A.: see Garrison, Erickson, Patten, and Timossi.
- Fitzgerald, C.S., P.R. Krausman, and M.L. Morrison: Use of buried and non-buried traps to sample desert rodents, 140-143.
- Foster, M.S.: see Schaeffer, Foster, Landrau, and Walder.
- Garrison, B.A., R.A. Erickson, M.A. Patten, and I.C. Timossi: California Wildlife Habitat Relationships System: Effects of county attributes on prediction accuracy for bird species, 87-101.
- Garrison, B.A.: see Timossi, Garrison, and Barrett.
- Goldberg, S.R.: Reproduction in the red diamond rattlesnake in California, 177-180.
- Goldberg, S.R. and C.R. Bursey: First reported occurrence of *Ophidasacaris labiatopapillosa* (Nematoda: Ascarididae) in the red diamond rattlesnake, 181-182.
- Gonzalez-Acosta, A.F., G. De la Cruz-Agüero, R. Rodriguez-Sanchez, J. De la Cruz-Agüero, and J.L. Castro-Aguirre: First records of morphological abnormalities in leatherjack, 79-82.
- Gordus, A.G.: A blood fluke from a northern pintail in California, 85-86.

- Hawkins, S.W. and J.M. Tipping: Predation by juvenile hatchery salmonids on wild fall chinook salmon fry in the Lewis River, Washington, 124-129.
- Hoff, G.R.: Range extensions of 3 species of macrourids from the west coast of North America, 113-117.
- Jow, T., J. Baxter, and R. Klingbeil: In memoriam: Patricia Powell, 146.
- Klingbeil, R.: see Jow, Baxter, and Klingbeil.
- Kohlhorst, D.W.: Status of striped bass in the Sacramento-San Joaquin Estuary, 31-36.
- Kohlhorst, D.W.: see Schaffter and Kohlhorst.
- Krausman, P.R.: see Fitzgerald, Krausman, and Morrison.
- Landrau, M.E.: see Schaeffer, Foster, Landrau, and Walder.
- Laughlin, L. and J. Ugoretz: An observation of blue shark parturition in the Southern California Bight, 83-84.
- Lavenberg, R.J.: see Posner and Lavenberg.
- Liu, F.-G., S.-D. Yang, and H.-C. Chen: Effects of stocking density on growth, gross composition, and plasma and hepatic metabolite levels in palmetto bass, *Morone saxatilis* x *M. chrysops*, 11-20.
- Magee, A., C.A. Myrick, and J.J. Cech, Jr.: Thermal preference of female threespine sticklebacks under fed and food-deprived conditions, 102-112.
- Morrison, M.L.: see Fitzgerald, Krausman, and Morrison.
- Myrick, C.A.: see Magee, Myrick, and Cech.
- Ochoa-López, E.: see Salazar-Hermoso, Ochoa-López, and Villavicencio-Garayzar.
- Patten, M.A.: see Garrison, Erickson, Patten, and Timossi.
- Pauli, A.M.: see Bleich and Pauli.
- Pierce, B.M.: see Bleich and Pierce.
- Pondella, D.J., II: First record of deepwater bass, *Serranus aequidens* (Serranidae), from California, 130-134.
- Posner, M. and R.J. Lavenberg: Movement of California halibut along the coast of California, 45-55.
- Rodriguez-Sanchez, R.: see Gonzalez-Acosta, Cruz-Agüero, Rodriguez-Sanchez, Cruz-Agüero, and Castro-Aguirre.
- Rosales-Casián, J.A. and G. Ruiz-Campos: Northern range extension of the white grunt, *Haemulopsis leuciscus*, 135-137.
- Ruiz-Campos, G.: see Rosales-Casián and Ruiz-Campos.
- Salazar-Hermoso, F., E. Ochoa-López, and C. Villavicencio-Garayzar: Stranding records of the oarfish in and around Bahía de la Paz, Mexico, 70-74.
- Schaefer, K.M. and J. Childers: Northernmost occurrence of the slender tuna, *Allothunnus fallai*, in the Pacific Ocean, 121-123.
- Schaefer, R.J.: Biological characteristics of mule deer in California's San Jacinto Mountains, 1-10.
- Schaeffer, T.N., M.S. Foster, M.E. Landrau, and R.K. Walder: Diver disturbance in kelp forests, 170-176.
- Schaffter, R.G. and D.W. Kohlhorst: Status of white sturgeon in the Sacramento-San Joaquin Estuary, 37-41.
- Seigel, J.A.: see Ugoretz and Seigel.
- Sweetnam, D.A.: Status of delta smelt in the Sacramento-San Joaquin Estuary, 22-27.
- Timossi, C., B.A. Garrison, and R.H. Barrett: In memoriam: Irene Timossi, 183.
- Timossi, I.C.: see Garrison, Erickson, Patten, and Timossi.
- Tipping, J.M.: see Hawkins and Tipping.
- Ugoretz, J.: see Laughlin and Ugoretz.

- Ugoretz, J.K. and J.A. Seigel: First eastern Pacific record of the goblin shark, *Mitsukurina owstoni* (Lamniformes: Mitsukurinidae), 118-120.
- Unitt, P.: Review of: The Marin County breeding bird atlas and atlas of the breeding birds of Monterey County, California, 42-43.
- Villavicencio-Garayzar, C.: see Salazar-Hermoso, Ochoa-López, and Villavicencio-Garayzar.
- Walder, R.K.: see Schaeffer, Foster, Landrau, and Walder.
- Yang, S.-D.: see Liu, Yang, and Chen.
- Young, J.N.: see Bettaso and Young.

## SUBJECT

- Abnormalities, morphological: in leatherjack, 79-82.
- Ascarid: occurrence in red diamond rattlesnake, 181-182.
- Attributes, county: effect on prediction accuracy for bird species by the California Wildlife Habitat Relationships System, 87-101.
- Bass,  
  deepwater: first record from California, 130-134  
  palmetto: effects of stocking density, 11-20.  
  striped: status in the Sacramento-San Joaquin Estuary, 31-36.
- Characteristics, biological: mule deer in the San Jacinto Mountains, 1-10.
- Collars, long-term radio: juvenile mule deer, 56-62.
- Colorado River, lower: striped mullet and Pacific tenpounder, 85-76.
- Composition, gross: effect of stocking density in palmetto bass, 11-20.
- Deer,  
  black-tailed: movement across Trinity Reservoir, 63-69.  
  mule:  
    biological characteristics in the San Jacinto Mountains, 1-10.  
    long-term radio collars, 56-62.
- Density, stocking: effects on palmetto bass, 11-20.
- Diver disturbance: kelp forests, 170-176.
- East Mojave National Scenic Area: hunting and trapping, 148-160.
- Estuary, Sacramento-San Joaquin:  
  status of delta smelt, 11-27.  
  status of striped bass, 31-36.  
  status of white sturgeon, 37-41.
- Fluke, blood: in northern pintail, 85-86.
- Growth: effect of stocking density in palmetto bass, 11-20.
- Grunt, white: northern range extension, 135-137.
- Habitats, shallow-water and channel: use by delta smelt, 161-169.
- Halibut, California: movement along the California coast, 45-55.
- Hunting: East Mojave National Scenic Area, 148-160.
- Kelp forests: diver disturbance, 170-176.
- Leatherjack: morphological abnormalities, 79-82.
- Macrourids: range extensions of 3 species, 113-117.
- Metabolite levels, plasma and hepatic: effect of stocking density in palmetto bass, 11-20.
- Movement:  
  black-tailed deer across Trinity Reservoir, 63-69.  
  California halibut, 45-55.

Mullet, striped: spawning in the lower Colorado River, 75-76.

Oarfish: stranding in and around Bahia de la Paz, Mexico, 70-74.

Parasitism:

*Ophidascaris labiatopapillosa* in red diamond rattlesnake, 181-182.

Blood fluke in northern pintail, 85-86.

Parturition: blue shark in the Southern California Bight, 83-84.

Pintail, northern: blood fluke, 85-86.

Predation: juvenile hatchery salmonids on wild fall chinook salmon fry, 124-129.

Prediction accuracy: effect of county attributes using the California Wildlife Habitat Relationships System, 87-101.

Range extension:

chinook salmon in Mexico, 77-78.

deepwater bass, 130-134.

goblin shark, 118-120.

macrourids, 113-117.

Pacific tenpounder, 75-76.

slender tuna, 121-123.

white grunt, 135-137.

Rattlesnake, red diamond:

occurrence of *Ophidascaris labiatopapillosa*, 181-182.

reproduction, 177-180.

Reproduction: red diamond rattlesnake, 177-180.

Review, book:

Avian conservation. Research and management, 144-145.

Marin County breeding bird atlas, 42-43.

Atlas of the breeding birds of Monterey County, 42-43.

Rodents, desert: buried and non-buried traps, 140-143.

Salmon, chinook:

first Mexican record, 77-78.

predation by hatchery salmonids, 124-129.

Salmonids, juvenile hatchery: predation on wild fall chinook salmon, 124-129.

San Jacinto Mountains: biological characteristics of mule deer, 1-10.

Shark,

blue: observation of parturition, 83-84.

goblin: first eastern Pacific record, 118-120.

Smelt, delta:

juvenile use of shallow water and channels, 161-169.

status in the Sacramento-San Joaquin Estuary, 23-27.

Spawning: by striped mullet in the lower Colorado River, 75-76.

Splittail: status in California, 28-30.

Sticklebacks, threespine: thermal preference, 102-112.

Stranding: oarfish in Bahia de la Paz, Mexico, 70-74.

Sturgeon, white: status in the Sacramento-San Joaquin Estuary, 37-41.

Survivorship: black-tailed deer across Trinity Reservoir, 63-69.

Tenpounder, Pacific: return to the lower Colorado River, 75-76.

Thermal preference: threespine sticklebacks, 102-112.

Track: casting method, 138-139.

Trapping: East Mojave National Scenic Area, 148-160.

Traps, buried and non-buried: desert rodents, 140-143.

Trinity Reservoir: movement and survivorship of black-tailed deer, 63-69.

Tuna, slender: northernmost occurrence in the Pacific Ocean, 121-123.

Wildlife Habitat Relationships System, California: prediction accuracy for bird species, 87-101.

## SCIENTIFIC NAMES

- Acacia constricta*, 141.  
*Acipenser medirostris*, 37.  
*Acipenser transmontanus*, 37-41.  
*Adenostoma fasciculatum*, 2.  
*Agkistrodon piscivorus*, 177.  
*Alectoris chukar*, 154.  
*Allothunnus fallai*, 121-123.  
*Amphiura tridactylum*, 182.  
*Anas acuta*, 85-86.  
*Anisus vortex*, 85.  
*Anthopleura* spp., 173.  
*Arctostaphylos* spp., 2.  
*Artemesia tridentata*, 57.  
*Artemia* sp., 104.  
*Asterina miniata*, 173.  
*Astropecten* spp., 173.  
*Aythya valisineria*, 85.  
*Baiomys taylori*, 142.  
*Calliostoma* spp., 173.  
*Cancer* spp., 173.  
*Ceanothus greggii*, 57.  
*Ceanothus* spp., 2.  
*Cercidium*, spp., 2.  
*Cercocarpus betuloides*, 57.  
*Cercocarpus ledifolius*, 2.  
*Cervus elaphus*, 56.  
*Chaetodipus baileyi*, 141.  
*Chaetodipus penicillatus*, 142.  
*Chondracanthus* spp., 173.  
*Chrysothamnus nauseosum*, 57.  
*Chrysothamnus* spp., 2.  
*Coleogyne ramosissima*, 57.  
*Coryphaenoides cinereus*, 113-17.  
*Cottus extensus*, 109.  
*Coluber constrictor*, 181.  
*Coluber constrictor flaviventris*, 182.  
*Crotalus molossus*, 179.  
*Crotalus ruber*, 177-180, 181-182.  
*Crotalus tigris*, 179.  
*Cymatogaster aggregata*, 108.  
*Dendritobilharzia pulverulenta*, 85-86.  
*Dermacentor albipictus*, 6.  
*Dermacentor hunteri*, 6, 8.  
*Dermacentor occidentalis*, 6.  
*Dipodomys* spp., 140.  
*Elops affinis*, 76.  
*Engraulis mordax*, 80, 162.  
*Ephedra* spp., 57.  
*Felis concolor*, 68.  
*Gasterosteus aculeatus*, 102-112.  
*Haemulopsis leuciscus*, 135-137.  
*Henrichia* spp., 173.  
*Heterodon platirhinus*, 181.  
*Hilaria mutica*, 141.  
*Hypomesus nipponensis*, 23.  
*Hypomesus olidus*, 23.  
*Hypomesus transpacificus*, 22-27, 33, 161-169.  
*Istiophorus platypterus*, 121.  
*Lampropeltis getula*, 181.  
*Larrea tridentata*, 2, 141.  
*Leiostomus xanthurus*, 167.  
*Lepus californicus*, 154.  
*Leuresthes tenuis*, 108.  
*Linognathus ovillus*, 5.  
*Lophortyx gambelii*, 152.  
*Macrocystis pyrifera*, 170-176.  
*Malacocephalus laevis*, 113-117.  
*Masticophis taeniatus*, 177.  
*Mazzaella* spp., 173.  
*Mesocetoides* sp., 181.  
*Mitsukurina owstoni*, 118-120.  
*Morone saxatilis*, 31-36.  
*Morone saxatilis* x *M. chrysops*, 11-20.  
*Mugil cephalus*, 75-76.  
*Neolipoptena ferrisi*, 6.  
*Neotoma albigula*, 141.  
*Nerodia cyclopion*, 181.  
*Nerodia erythrogaster*, 181.  
*Nerodia rhombifer*, 182.  
*Nerodia sipedon*, 182.  
*Nezumia tirolepis*, 113-117.  
*Odocoileus hemionus*, 56-62, 68, 152.  
*Odocoileus hemionus californicus*, 2.

- Odocoileus hemionus columbianus*, 63-69.  
*Odocoileus hemionus fuliginatus*, 1-10.  
*Odocoileus hemionus hemionus*, 7.  
*Odocoileus virginianus*, 7.  
*Oligoplites saurus*, 79-82.  
*Olivella biplicata*, 173, 174.  
*Oncorhynchus chrysogaster*, 77.  
*Oncorhynchus clarki*, 124-129.  
*Oncorhynchus gorboscha*, 125.  
*Oncorhynchus keta*, 125, 167.  
*Oncorhynchus kisutch*, 124-129.  
*Oncorhynchus mykiss*, 77, 108, 124-129.  
*Oncorhynchus mykiss whitei*, 109.  
*Oncorhynchus nerka*, 125.  
*Oncorhynchus tshawytscha*, 33, 77-78, 124-129, 167.  
*Onychomys* spp., 141.  
*Ophidascaris labiatopapillosa*, 181-182.  
*Oregonia gracilis*, 173.  
*Orthasterias koehleri*, 173.  
*Ovis canadensis*, 61, 152.  
*Ovis canadensis cremnobates*, 2.  
*Ovis canadensis nelsoni*, 2.  
*Ovis musimon*, 56.  
*Pachycerianthus fimbriatus*, 173.  
*Paralabrax auroguttatus*, 130.  
*Paralabrax clathratus*, 130.  
*Paralabrax maculatofasciatus*, 130.  
*Paralabrax nebulifer*, 130.  
*Paralichthys californicus*, 45-55.  
*Paralichthys* spp., 167.  
*Peromyscus* spp., 142.  
*Phoca vitulina*, 173.  
*Phyllospadix* spp., 173.  
*Pinus jefferi*, 2.  
*Pinus ponderosa*, 2.  
*Pisaster* spp., 173.  
*Pituophis catenifer*, 177.  
*Planorbis planorbis*, 85.  
*Plecoglossus altivelis*, 17.  
*Pogonichthys macrolepidotus*, 28-30.  
*Pugettia* spp., 173.  
*Prionace glauca*, 83-84.  
*Prosopis velutina*, 141.  
*Prunus andersonii*, 57.  
*Purshia tridentata*, 57.  
*Quercus dumosa*, 2.  
*Rana capito*, 182.  
*Rana catesbeiana*, 182.  
*Rana sphenoccephala*, 182.  
*Regalecus glesne*, 70-74.  
*Regalecus kanoi*, 70-74.  
*Regalecus russelli*, 70-74.  
*Reithrodontomys* spp., 142.  
*Salmo salar*, 108.  
*Salvelinus alpinus*, 11.  
*Salvelinus fontinalis*, 108.  
*Serranus aequidens*, 130-134.  
*Serranus fasciatus*, 130.  
*Serranus huascarii*, 130, 131.  
*Sigmodon* spp., 141.  
*Sylvilagus audubonii*, 154.  
*Tegula* spp., 173.  
*Theragra chalcogramma*, 108.  
*Thunnus alalunga*, 121.  
*Zalophus californianus*, 131, 173.  
*Zenaidura macroura*, 154.  
*Zostera marina*, 135.  
*Zostera* sp., 104.

